Increase in quarterly HIV testing and UAI amongst MSM in London with decrease in anxiety

Updates from a continuous online assessment tool

Wang, D; Dunkley, Y; West, R; Skipper, K; Worrall, S; Evans, A; Ussher, G; Santos, M



Positive Eas

Outline



Who are we



What was the online questionnaire about



Findings from the questionnaire



What to expect next









Who are we?

Over the last 10 years, we have:

- Delivered over 4,200 sessions in more than 150 venues and key events (e.g. Pride and National HIV testing week)
- ❖ Engaged with more than 428,000 people during the outreach sessions
- ❖ Talked to over 206,000 men regarding HIV and sexual wellbeing related issues
- ❖ BASK assessed more than 17,000 MSMs (face to face)

Between 2008 and 2014,

- * 4,731 counselling sessions were provided to 661 MSM
- * 1,445 mentoring sessions were provided to 253 MSM

Since April 2013, we have:

conducted about 5,000 HIV tests





018 HIV Prevention Survey for Gay

Before hitting the "Submit" button at the end of this survey, you can find further information an

HIV Testing

If you are sexually active, it is recommended that you test for HIV and STIs at least once a year, reparalless of your shaeton, and more frequently if you are having unprotected sex will casual partners. For more information on HIV starting, self-heating, path-sampling, where to sext and HIV transmission visit tha Do it London website, www.doltondon.org/hiv/sating

Condoms

Listing a condom during set is all the best way to protect yourself against HII and account bearsembled infections, LEPP fined beardors Shop to desire fine lettle quality modernal and label to varies across London. You can pill-king your fine condoms and label in over 50 gay versues and 40 AIM Personantils contracts service. You can also protesse a huge variety or lamosat condoms and bleas via the President and those website. Please visit www.dolfstondom.orbitoschers for more information.

Pre-exposure prophylaxis (PrEP)

PYEP is a HIV prevention strategy that would use entiretrovirals (ARIVs) to protect HIVnegative people from HIV infection, in this strategy, people would take the medications before they were exposed to HIV to lower their risk of infection. For more information on PYEP visit www.dottoncom.oru/PYEP.

Post-exposure prophylaxis (PEP)

If you have had unprotected sex (including condom failure incidents) or shared drug injecting equipment with somebody who might be HIV positive, then you may consider PE a one-month course of HIV treatment intended to reduce the risk of you becoming infecte with HIV.

To have the best chance of it being effective you need to start taking PEP as soon possible after the possible exposure to HIV and certainly within 72 hours. For more information on PEP (volt the NAM website https://www.aidamag.com/Possions.edu/

Undetectable

Pagos in ring with HIV on ART justnesshorks having of with an undisticable virial load in their blood hear an eightlight size of season transmission of HIV. Depending on the virial amplity of may take as ting as any mortes for the virial load to become undisticable. Contributed and listilities HIV appreciation registers selection of appropriate significant scalables arbeitness to treatment. HIV virial suppression should be monitored to assure both enterous lineatine and untilin based hearing.

NOTE in unpatcable viril load is hybidally under All popelarin depending on the dispondin sets, however, shalles below a person line, with Viril an entirely interest because (ART) with a viril load of 200 operation or less also cannot sexually treated HM. This is a state being "Viril yopposses of the Unit proposed of the Unit parent plan and any plan and the Unit parent plan and any operation and any op

For more information on U+U visit <u>www.doitiondon.org/Undetectabl</u>

Further Support


If you are taking risks and sar worked about it, get in touch with us at DMI and we can takly you through all the support options in your area. We can legi you information on ebbasiliochange services such as individual and group support, link you in with specialist LISET on and alchord services and provide information or your local IMES services. You can either email DMI - info@pringerheasitio.org.uji or visit the DMI website for further




The Assessment Tool – 14Q


- BASK to 14Q under LHPP contract (Do it London)
- From July 2015, we started using online questionnaires selfassessment toll: http://www.gmipartnership.org.uk/gmi-survey/
 - Target: MSM, dating app users
 - Aims:
 - To assess behavioural risks, e.g. UAI and chemsex
 - To promote up-to-date HIV prevention information, e.g. HIV testing, PrEP, PEP, and Undetectable equals Untransmittable (U=U)
 - Observe attitudes and collect thoughts regarding the prevention methods
 - To engage with those who might need further support
- Primarily broadcast via Grindr
- Evolving responding to DiL campaigns and trends within the community

Findings - general





• 12,250 responses since July 2015






gmi partnership






Findings -trends

Compare Q4 data over 4 years,



*Among those who have had UAI with casual partners,



Finding - limitation



LIMITS OF SAMPLING METHOD



LIMITS OF QUESTIONS



LIMITS IN INTERPRETATIONS







Continue and modify Q14

Secure provisions

Identify key information and key questions



Research partner

Academic institutes
Research professionals



In-depth research

Quantitative & qualitative
Reasons behind changes of
behaviours and attitudes
Specific target groups, e.g. over
50s.





info@gmipartnership.org.uk

Special thanks to:

- GMI outreach team: Failla, G; Gillard, N; Mosaheb, Y; Nardi, M; Wragg, P
- GMI communication lead: Delacour, M; Neilson, A;
- Do it London

